



Challenges in Genome-Wide Location Analysis



The ChIP-DASL Technology



5

Q

 ∞

9


5


Specificity and Sensitivity of DASL

;											
Kector #	-	7	8	エ	5	5 6 7	14	α	<u>a</u>	0	
Genomic DNA		105	105	105	105 105 105 105	105				105	
Spiked Plasmid DNA			102	103	104	102 103 104 105 105	105		105		
Oligo Pool for Genomic DNA								+	+	+	
Oligo Pool for Spiked DNA	+	+	+	+	+	+	+				
										A Party Commence	
											l diam


F16.34

Characterization of the ChIP-DASL Technology by using Spiking Controls







5-fold Spiking on the 2K Human Promoter Array



Identification of Estrogen Receptor Target Genes by DASL




F16, 6




Mapping Transcription Units by Tiling on the β-globin Locus

Project goal: Mapping functional DNA elements in the ENCODE region Collaborator: Bing Ren (UCSD)



Additional Applications of DASL



This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
□ OTHER:

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.